

Validity of hydrostatic balance in the perturbations generated by the Met Office high resolution ensemble forecast system

Sanita Vetra, S. Migliorini, M. Dixon, N. K. Nichols, S. Ballard

May 20, 2009

Funding: NERC & Met Office

Motivation

Data assimilation algorithm:

$$x^a = x^f + K(y - Hx^f)$$

$$K = PH^T(HPH^T + R)^{-1}$$

- ▶ Initialization of P is very important especially for variational methods
- ▶ Incorrect specification of forecast errors can lead to a degraded forecast


The questions:

- ▶ Does hydrostatic balance holds in forecast perturbations at high resolutions?
- ▶ If it does then to what extent?

Case study of 3h window on 27/07/2008 over southern UK:

- ▶ Model used is Met Office Global and Regional Ensemble Prediction System (MOGREPS)
- ▶ Model has 1.5km horizontal resolution
- ▶ MOGREPS uses ETKF methodology to determine a set of perturbed forecasts and 1.5km UM forward model
- ▶ Model is initialized with a reconfigured ensemble from 24km to 1.5km at 18Z
- ▶ 24 ensemble member forecasts available at 19Z, 20Z, 21Z
- ▶ Domain size $360 \times 288 \times 70$

- └ Case study
- └ Data selection



Available fields for all vertical levels are

- ▶ Exner pressure Π
- ▶ potential temperature θ
- ▶ and specific humidity q

Let the vertical state vector be defined as

$$\mathbf{x} = (\Pi_{0,\dots,k-1}, \theta_{0,\dots,k-1}, q_{0,\dots,k-1}),$$

where $k = 70$ is number of vertical levels and an ensemble is defined as

$$\mathbf{X} = [\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{N-1}] \in \mathcal{R}^{3k \times N}$$

where $N = 24$ is the number of ensemble members.

The ensemble mean is defined as $\bar{\mathbf{x}}$. Thus, the ensemble perturbations are given by

$$\mathbf{x}'_i = \mathbf{x}_i - \bar{\mathbf{x}}$$

where $i = 0, \dots, N - 1$.

Hydrostatic balance is given by (e.g. see Wallace 2006),

$$\frac{dp}{dz} = -\frac{gp}{RT}$$

where $p(z)$ is pressure, $T(z)$ is temperature, z is vertical height, and $R = 287.06 \text{ J K}^{-1}\text{kg}^{-1}$ is the gas constant for dry air.

Using expression for Exner pressure $\Pi = \left(\frac{p}{p_0}\right)^{R/c_p} = \frac{T}{\theta^v}$ and virtual potential temperature $\theta^v = \theta (1 + (\epsilon^{-1} - 1)q)$, hydrostatic equation above can be written as follows

$$\frac{d\Pi}{dz} = -\frac{g}{c_p} (1 + (\epsilon^{-1} - 1)q)^{-1} \theta^{-1}.$$

By decomposing $\Pi = \bar{\Pi} + \Pi'$, $\theta = \bar{\theta} + \theta'$ and $q = \bar{q} + q'$ above equation may be linearized giving a first order approximation to hydrostatic equation for the perturbations:


$$\frac{d\Pi'}{dz} = \frac{g}{c_p} \left[\frac{(\epsilon^{-1} - 1)q'}{(1 + (\epsilon^{-1} - 1)\bar{q})^2\bar{\theta}} + \frac{\theta'}{(1 + (\epsilon^{-1} - 1)\bar{q})\bar{\theta}^2} \right].$$

Rearranging gives

$$\theta'_H = -\frac{(\epsilon^{-1} - 1)q'\bar{\theta}}{1 + (\epsilon^{-1} - 1)\bar{q}} + \frac{c_p}{g} \frac{d\Pi'}{dz} \bar{\theta}^2 (1 + (\epsilon^{-1} - 1)\bar{q}).$$


└ Results

└ Perturbation correlation




- ▶ Compute θ'_H for each column
- ▶ Compare correlation matrices of θ' and θ'_H
- ▶ Examine explained variances and error flow
- ▶ Aggregate data around Conv19Z and NonConv points to examine hydrostatic balance as a function of horizontal scale

Correlation matrices for 19Z at 1.5km resolution




(a) θ' , Non-Conv (b) θ'_H , Non-Conv




(c) θ' , Conv19Z (d) θ'_H , Conv19Z


Correlation matrices for 19Z at 4km and 12km resolution




(a) θ' , Conv4km



(b) θ'_H , Conv4km



(c) θ' , Conv12km



(d) θ'_H , Conv12km


Balance measure

The explained variance is given by


$$E(z) = \left(1 - \frac{\sigma_U^2(z)}{\sigma^2(z)} \right)$$

where σ^2 is the grid-point variance of θ' and σ_U^2 is the variance of the unbalanced part of the perturbations, i.e. $\theta'_U = \theta' - \theta'_H$, and z is the vertical level. Thus, if $E \approx 1$ then perturbations are close to hydrostatic balance and if $E \approx 0$ then they are imbalanced.


Explained variance at 1.5km resolution



(e) $E(z)$ at 19Z




(f) $E(z)$ at 20Z




(g) $E(z)$ at 21Z

Explained variance at coarser resolution



$$rel.error = \frac{\sqrt{(\theta'_H - \theta')^2}}{|\theta'_H|} \times 100$$



(a) Non-Conv

(b) Conv19Z

(c) Conv20Z

(d) Conv21Z

- ▶ At 1.5km resolution hydrostatic balance does not hold in the perturbations in the regions of convection but it does hold in the regions where convection is not present.
- ▶ This suggests that hydrostatic balance should be relaxed around convective columns in the correlation matrices at 1.5km resolution. A way of achieving this would be by redesigning the control variable transform in UM.
- ▶ $\approx 20\text{km}$ horizontal resolution is the limit at which the hydrostatic balance becomes valid over the entire domain.

Future work

- ▶ Investigate balance properties in P using an idealised 1+2D convective model and EnSRF
- ▶ Further, how balances are affected by applying localization
- ▶ Test performance of EnSRF when applied to a convective model

Thank You!






Fig. 7: Setup and flow of the MORGREPS ensemble prediction system. Here $t_0 = 18Z$, $t_1 = 19Z$, $t_2 = 20Z$, and $t_3 = 21Z$ all on 27/07/2008 available at (**)



(a) 19Z

(b) 20Z



(c) 21Z