

Influence of mid-latitude cyclones on European background surface ozone investigated in observations, MACC and MERRA-2 reanalyses

K. Emma Knowland^{1,2}, Ruth Doherty³, Kevin Hodges⁴ and Lesley Ott²

¹USRA/GESTAR; ²NASA GSFC, Global Modeling and Assimilation Office (GMAO), Greenbelt, MD;

³School of Geoscience, University of Edinburgh, Edinburgh, UK; ⁴Department of Meteorology, University of Reading, Reading, UK



Take Home Messages!

First study to our knowledge to quantify the influence extratropical cyclones have on the temporal variability of springtime surface ozone (O_3) measured on the west coast of Europe when cyclones are nearby.

We show **passing cyclones have a discernible influence on surface O_3 concentrations.**

In-depth findings from four case studies, using a combination of reanalyses and a modeled tracer, demonstrate there are several transport pathways before O_3 -rich air eventually reaches the surface. (Knowland et al., 2017 ACP)

Surface O_3 Observations

Location	Period	Source	Notes
Mace Head, Ireland	1988-2012	EMEP	In main storm track (Fig. 1)
Monte Velho, Portugal	1989-2009*	EMEP	South of main storm track

Reanalysis

Reanalysis	Period	Model	O_3 Chemistry	O_3 Assimilated	Resolution	Reference
ERA-Interim	1979-present	ECMWF IFS	O_3 chemistry parametrization	O_3 assimilated	~0.7° 60 levels	Dee et al., 2011
MACC	2003-2012	ECMWF IFS	Coupled to MOZART-3 CTM	O_3 assimilated	~0.7° 60 levels	Inness et al., 2013
MERRA-2	1980-present	GEOS-5	Simplified O_3 chemistry	O_3 assimilated	~0.5° 72 levels	Gelaro et al., 2017

- Met and chemical variables on 12 pressure levels from 1000 to 200 hPa
- Relative vorticity at 850 hPa (ζ_{850hPa}), mean sea level pressure (MSLP), temperature (T), specific humidity (q), winds (u,v), vertical velocity (ω), equivalent potential temperature (θ_e), and O_3 .




Figure 1: Storm track density (contours) over Europe, North, Center and South regions indicated for Mace Head (blue dot), Monte Velho (red dot).


Storm tracks and O_3

- Storm tracks were identified in ERA-Interim and MACC using the objective feature tracking algorithm, TRACK (Hodges 1995, 1999).
- O_3 at Mace Head and Monte Velho were sorted each season, to remove the increasing background signal, and ranked by percentiles (pc).
- Tracks were matched to concurrent surface O_3 observations at Mace Head and Monte Velho.

Region	% tracks "high" O_3 $O_3 > 75^{th}$ pc	# years more tracks with high O_3 (# significant)	% tracks "low" O_3 $O_3 > 75^{th}$ pc	# years more tracks with low O_3 (# significant)
North	52 %	18 (15)	37 %	5 (0)
Center	51 %	17 (6)	41 %	6 (1)
South	45 %	7 (2)	53 %	16 (8)

When cyclones track north of 53°N, there is a significant relationship with high levels of surface O_3 ($> 75^{th}$ pc). The further away a cyclone is from the main storm track, more likely associated with low O_3 ($< 25^{th}$ pc).

- Case study cyclones (Fig 2 below) identified for
 - high O_3
 - Passing through North or South regions
 - Strong, top 20 % based on maximum ζ_{850hPa}
 - two consecutive time steps with high O_3



Strong cyclone tracks, N1 and S1, associated with high O_3 at Mace Head (Fig. 2)

- All points within a 20° spherical cap over the storm centre are selected and oriented in the direction of the storm's movement




Figure 3: Schematic of a storm track identified on 850hPa field, transferred on multiple levels. Spherical cap around storm center at time of maximum ζ_{850hPa} is illustrated.

N1 cyclone: North of Mace Head




Figure 4: N1 cyclone is a secondary cyclone to parent low "P". a) 1000 hPa MACC O_3 , SLP, winds. b) Analysis chart from UK Met Office. Mace Head indicated by pink circle.

Strong O_3 gradient across the cold front, 1) with maritime clean air first to Mace Head 2) elevated levels of O_3 behind the front.




Figure 5: N1 cyclone 4 March 2007 00 UTC




Figure 6: N1 cyclone 24 hours later (5 March 2007 UTC). Cold front has passed over Mace Head.

Descent of O_3 -rich air behind the strong frontal boundary in dry intrusion (DI) airstream.
Ascent of O_3 -poor air in the warm conveyor belt (WCB) up to 400 hPa.
Descent of O_3 -rich air in DI
Transport across North Atlantic in strong westerly winds.
Persistent high O_3 observed at Mace Head for 1.5 days

S1 cyclone: South of Mace Head




Figure 5: S1 cyclone 25 April 2012 00UTC in MACC




Figure 5: S1 cyclone 25 April 2012 00UTC MERRA-2

- Tropopause is level above Mace Head. DI is to the south.
- O_3 -rich air below tropopause in MERRA-2 -> air is likely from the stratosphere
- Residual high O_3 from upstream decaying DI airstream entrained into S1

Figure key: O_3 (color), tropopause (2 PVU, thick black line), ω (white=descent, black=ascent), MSLP (lowest pressure level), θ_e (dotted lines), Mace Head (black dot)

Dee et al., 2011 "The ERA-Interim reanalysis: configuration and performance of the data assimilation system" QJ.R. Meteorol. Soc., 137, 553-597. doi:10.1002/qj.828.
Gelaro et al., 2017 "The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2)" J. Climate, 30, 5419-5454. doi:10.1175/JCLI-D-16-0758.1.
Hodges, K.I. 1995 "Feature tracking on the unit-sphere" Mon. Wea. Rev., 123, 3458-3465.
Hodges, K.I. 1999 "Adaptive constraints for feature tracking" Mon. Wea. Rev., 127, 1362-1373
Inness et al., 2013 "The MACC reanalysis: an 8 yr data set of atmospheric composition" Atmos. Chem. Phys., 13, 4073-4109
Knowland et al., 2017 "The influence of mid-latitude cyclones on European background surface ozone", Atmos. Chem. Phys., 17, 12421-12447. doi:10.5194/acp-17-12421-2017